

A close-up photograph of a woman with dark hair, smiling warmly and looking towards a young child. The child is wearing a yellow shirt and looking back at the woman. The background is a plain, light-colored wall.

**“A good day seems
full of possibility”**

— Diana, mom of Lina

ZTALMY® could mean more days with fewer seizures for your loved one with CDKL5 deficiency disorder (CDD).

Lina is living with CDD.

What is ZTALMY?

- ZTALMY is a prescription medicine used to treat seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD) in people 2 years of age and older.
- ZTALMY is a federally controlled substance (CV) because it contains ganaxolone that can be abused and lead to dependence. Keep ZTALMY in a safe place to prevent misuse and abuse. Selling or giving away ZTALMY may harm others and is against the law. Tell your healthcare provider if you or your child have ever abused or been dependent on alcohol, prescription medicines or street drugs.
- It is not known if ZTALMY is safe and effective in children under 2 years of age.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

The logo for Ztalmy, featuring the brand name in a stylized purple font with a registered trademark symbol. Below it, the generic name (ganaxolone) and formulation (oral suspension) are listed, along with the concentration (50 mg/mL).

Ztalmy®
(ganaxolone) oral suspension | 50 mg/mL

Seizures in CDD

What would fewer seizures mean for your loved one with CDD?

Your family is likely familiar with how hard it is to control seizures associated with CDD. Your loved one may have tried several seizure medicines or may be on multiple medicines and still experiencing frequent seizures.

Seizures can disrupt daily life. They may cause your loved one to miss school, therapy, and other activities. Seizures can affect your family's balance of good days with CDD—the days with fewer seizures when you can interact more easily with your loved one or complete a therapy session.

Visit [ZTALMY.com/stories](https://www.ztalmymed.com/stories) to hear from families living with CDD

Important Safety Information

What is the most important information I should know about ZTALMY?

ZTALMY can cause serious side effects, including:

Sleepiness: ZTALMY may cause sleepiness. Taking ZTALMY with central nervous system (CNS) depressants including alcohol may increase sleepiness. Do not drive, operate heavy machinery, or do other dangerous activities until you know how ZTALMY affects you or your child.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

“

A bad day is always the same—holding Lina, rocking her, trying to soothe her.”

— Diana, mom of Lina

Lina is living with CDD.

How ZTALMY may help your loved one with CDD

↓ **31%**
SEIZURE
REDUCTION
with ZTALMY

vs

↓ **7%**
SEIZURE
REDUCTION
with PLACEBO

Proven effective in people 2 years and older with CDD

In the clinical study, ZTALMY significantly reduced the monthly seizure frequency over the 17-week treatment period. About half of participants (49) took ZTALMY and about half (51) received placebo. Those taking ZTALMY experienced a **4X reduction in median seizure frequency (31%), compared with placebo (7%).**

Reduced the frequency of seizures by at least half for some study participants

One in 4 participants taking ZTALMY had at least a 50% reduction in frequency of seizures vs 1 in 10 participants taking placebo.


1 in 4 experienced
at least a **50%**
seizure reduction

Individual results may vary.

Important Safety Information

Suicidal Thoughts or Actions: Like all other antiepileptic drugs, ZTALMY may cause suicidal thoughts or actions in a very small number of people, about 1 in 500. **Call your healthcare provider right away if you or your child have any of these symptoms, especially if they are new, worse, or worry you:**

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).



Lina is living with CDD.

Studied in people with treatment-resistant seizures associated with CDD

- 101 children and young adults with CDD aged 2 to 19 years participated in the clinical study
- Participants had previously tried and discontinued between 1 and 16 seizure medicines
- Participants were taking between 1 and 4 seizure medicines and were still having frequent seizures
- Seizure types that were assessed included bilateral tonic, generalized tonic-clonic, bilateral clonic, atonic, and focal to bilateral tonic-clonic

The first treatment indicated specifically for seizures associated with CDD in people 2 years and older

Important Safety Information (continued)

- thoughts about suicide or dying
- attempt to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Ztalmy®
(ganaxolone) oral suspension | 50 mg/mL

Side effects of ZTALMY

- In the clinical study, the most common side effects for ZTALMY (those that occurred in $\geq 5\%$ of participants and twice the rate of placebo) included*:
 - Sleepiness
 - Fever
 - Excessive salivation or drooling
 - Seasonal allergy
- 4% of participants taking ZTALMY discontinued treatment due to side effects vs 8% of participants taking placebo
 - Reasons for discontinuing ZTALMY were sleepiness and seizure (for 1 participant) and seizure (for 1 participant)

*These are not all of the possible side effects of ZTALMY. Talk to your healthcare provider for medical advice about side effects.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

In the clinical study, ZTALMY was used in combination with other seizure medicines

98% of participants taking ZTALMY were also taking other seizure medicines

Most common seizure medicines taken with ZTALMY: valproate, levetiracetam, clobazam, and vigabatrin.

Questions about ZTALMY? Visit [ZTALMY.com/resources](https://www.ztalmy.com/resources) to see FAQs

Important Safety Information

Pay attention to changes, especially sudden changes in mood, behaviors, thoughts, or feelings. Keep all follow-up visits with your healthcare provider as scheduled.

Stopping ZTALMY: Do not stop taking ZTALMY without first talking to your healthcare provider. Stopping ZTALMY suddenly can cause you or your child to have seizures more often or seizures that do not stop (status epilepticus).




Lina is living with CDD.

Ztalmy® 
(ganaxolone) oral suspension | 50 mg/mL

How ZTALMY works

ZTALMY is the first seizure medicine that is a neuroactive steroid. Exactly how ZTALMY works to treat seizures in CDD is not fully understood, but it is thought to reduce seizures by enhancing the activity of GABA (an inhibitory neurotransmitter) in the brain.



The role of GABA

- The exact cause of seizures in CDD, like in other epilepsies, is not fully known. Seizures may result from an imbalance of excitatory (stimulating) and inhibitory (calming) signals in the brain.
- GABA, the main inhibitory neurotransmitter, plays an important role in sending inhibitory signals to calm excited neurons (nerve cells). When these signals do not work as they should, neurons may become overexcited, which can lead to seizures.

Important Safety Information

What should I tell my healthcare provider?

Before taking ZTALMY, tell your healthcare provider about all of your or your child's medical conditions, including if you or your child:

- drink alcohol.
- have or have had depression, mood problems or suicidal thoughts or behavior.
- have abused or been dependent on prescription medicines, street drugs, or alcohol.
- have liver problems.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).



The ZTALMY difference

- GABA sends inhibitory signals by attaching to certain GABA receptors found on neurons. Depending on their location, some of these receptors produce short bursts of inhibition, while others produce continuous inhibition.
- ZTALMY is a different kind of seizure medicine that enhances both types of inhibition by working on certain GABA receptors in different locations on the neuron. This dual approach may help to calm overexcited neurons.

Important Safety Information (continued)

- are pregnant or plan to become pregnant. Tell your healthcare provider right away if you or your child become pregnant while taking ZTALMY. You and your healthcare provider will decide if you or your child should take ZTALMY while pregnant.
- are breastfeeding or plan to breastfeed. ZTALMY may pass into breast milk. Talk to your healthcare provider about the best way to feed your or your child's baby while taking ZTALMY.

Tell your healthcare provider about all the medicines you or your child take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ZTALMY may affect the way other medicines work, and other medicines may affect how ZTALMY works. Do not stop or start taking other medicines without talking to your healthcare provider.



What to know about taking ZTALMY

How should your loved one take ZTALMY?



Exactly as prescribed

3_x

3 times a day



With food

Before starting ZTALMY, read [How to Give ZTALMY](#) on ZTALMY.com.

Made with your loved one's needs in mind



Keto-friendly oral medicine.

ZTALMY is a sugar-free, cherry-flavored liquid, suitable for ketogenic diets.



Does not require refrigeration.

Keep ZTALMY in a safe place to prevent misuse and abuse.



ZTALMY dosing schedule

- For the first 28 days, your healthcare provider will likely increase your loved one's dose each week depending on how well your loved one is tolerating ZTALMY
- Give ZTALMY exactly as your healthcare provider tells you. Your healthcare provider will tell you how much ZTALMY to give and when to give it
- Use the bottle adaptor and oral dosing syringes provided by the specialty pharmacy to administer ZTALMY. Do not use a household teaspoon or tablespoon
- Your healthcare provider may adjust the dose of ZTALMY as your loved one's weight changes

Please see [Instructions for Use](#) for more information.

Important Safety Information

Especially tell your healthcare provider if you or your child take: alcohol; opioids; antidepressants.

Know the medicines you or your child take. Keep a list of them to show your healthcare provider or pharmacist when you or your child get a new medicine.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

Important Safety Information (continued)

What should be avoided while taking ZTALMY?

Do not drive, operate heavy machinery, or do other dangerous activities until you know how ZTALMY affects you or your child. ZTALMY may cause you or your child to feel sleepy.



Support for your ZTALMY® journey



Ztalmy one™

ZTALMY One is a comprehensive support program for you and your loved one.

ZTALMY One support includes:

- Benefits verification
- Prescription delivery
- ZTALMY cost support*
- Ongoing prescription support



ZTALMY One helps make treatment more *affordable*

Whether your family is insured, underinsured, or not insured at all, ZTALMY One has copay and prescription support programs that may help make treatment more affordable.

*Enrollment in ZTALMY One is required. Eligibility Terms and Conditions apply.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

Talking to your loved one's healthcare provider

Talk to your healthcare provider about the opportunity to reduce your loved one's seizures and what more "good days" could mean for your family. These questions can help you start a conversation about whether it is time to consider ZTALMY.



About seizure management in CDD

- Which seizure medicines have been studied for CDD?
- Are my loved one's seizure medicines effective at reducing the frequency of seizures associated with CDD?
- What is the goal of our seizure management plan?

About ZTALMY

- Can ZTALMY help us reach our seizure management goals?
- Will ZTALMY work with my loved one's current treatment plan?
- What are the potential risks and benefits of ZTALMY?

Bring these questions to your next appointment



Ztalmy®
(ganaxolone) oral suspension | 50 mg/mL

INDICATION

What is ZTALMY?

- ZTALMY is a prescription medicine used to treat seizures associated with cyclin-dependent kinase-like 5 (CDKL5) deficiency disorder (CDD) in people 2 years of age and older.
- ZTALMY is a federally controlled substance (CV) because it contains ganaxolone that can be abused and lead to dependence. Keep ZTALMY in a safe place to prevent misuse and abuse. Selling or giving away ZTALMY may harm others and is against the law. Tell your healthcare provider if you or your child have ever abused or been dependent on alcohol, prescription medicines or street drugs.
- It is not known if ZTALMY is safe and effective in children under 2 years of age.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about ZTALMY?

ZTALMY can cause serious side effects, including:

Sleepiness: ZTALMY may cause sleepiness. Taking ZTALMY with central nervous system (CNS) depressants including alcohol may increase sleepiness. Do not drive, operate heavy machinery, or do other dangerous activities until you know how ZTALMY affects you or your child.

Suicidal Thoughts or Actions: Like all other antiepileptic drugs, ZTALMY may cause suicidal thoughts or actions in a very small number of people, about 1 in 500.

Call your healthcare provider right away if you or your child have any of these symptoms, especially if they are new, worse, or worry you:

- thoughts about suicide or dying
- attempt to commit suicide
- new or worse depression
- new or worse anxiety
- feeling agitated or restless
- panic attacks
- trouble sleeping (insomnia)
- new or worse irritability
- acting aggressive, being angry, or violent
- acting on dangerous impulses
- an extreme increase in activity and talking (mania)
- other unusual changes in behavior or mood

Pay attention to changes, especially sudden changes in mood, behaviors, thoughts, or feelings. Keep all follow-up visits with your healthcare provider as scheduled.

Stopping ZTALMY: Do not stop taking ZTALMY without first talking to your healthcare provider. Stopping ZTALMY suddenly can cause you or your child to have seizures more often or seizures that do not stop (status epilepticus).

What should I tell my healthcare provider?

Before taking ZTALMY, tell your healthcare provider about all of your or your child's medical conditions, including if you or your child:

- drink alcohol.
- have or have had depression, mood problems or suicidal thoughts or behavior.

- have abused or been dependent on prescription medicines, street drugs, or alcohol.
- have liver problems.
- are pregnant or plan to become pregnant. Tell your healthcare provider right away if you or your child become pregnant while taking ZTALMY. You and your healthcare provider will decide if you or your child should take ZTALMY while pregnant.
- are breastfeeding or plan to breastfeed. ZTALMY may pass into breast milk. Talk to your healthcare provider about the best way to feed your or your child's baby while taking ZTALMY.

Tell your healthcare provider about all the medicines you or your child take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ZTALMY may affect the way other medicines work, and other medicines may affect how ZTALMY works. Do not stop or start taking other medicines without talking to your healthcare provider.

Especially tell your healthcare provider if you or your child take: alcohol; opioids; antidepressants.

Know the medicines you or your child take. Keep a list of them to show your healthcare provider or pharmacist when you or your child get a new medicine.

What should be avoided while taking ZTALMY?

Do not drive, operate heavy machinery, or do other dangerous activities until you know how ZTALMY affects you or your child. ZTALMY may cause you or your child to feel sleepy.

What are the side effects of ZTALMY?

See "What is the most important information I should know about ZTALMY?"

The most common side effects of ZTALMY include:

- sleepiness
- fever
- excessive saliva or drooling
- seasonal allergy

These are not all of the possible side effects of ZTALMY. For more information ask your healthcare provider or pharmacist. Tell your healthcare provider about any side effect that bothers you or your child or that does not go away. **Call your doctor for medical advice about side effects. You may report side effects to the FDA at www.fda.gov/medwatch or 1-800-FDA-1088.** You may also contact Immedica at 1-844-627-4687.

Please see [Full Prescribing Information](#) and [Medication Guide](#) or visit www.ztalmy.com for more information.





“
When it's a good day,
we can go outside or
we can play or we can do
some therapies together.”

— Diana, mom of Lina

Ask your healthcare provider
about ZTALMY and if it is right
for your loved one.

Visit [ZTALMY.com](https://www.ztalmy.com) for resources
and additional information.

Lina is living with CDD.

Please see additional Important Safety Information throughout and click here for [Medication Guide](#).

Marketed by:
Immedica Pharma US Inc., Chicago, IL 60642

©2026 Immedica Pharma US Inc. All rights reserved.
ZTALMY is a registered trademark of Immedica Pharma US Inc.
PRC-US-Ztalmy-00047-v4 1/26

